ELSEVIER

Environmental Pollution 146 (2007) I–VII

ENVIRONMENTAL POLLUTION

www.elsevier.com/locate/envpol

Contents of Volume 146

Number 1 2007

New Initiatives

1 **High uptake of 2,4,6-trinitrotoluene by vetiver grass – Potential for phytoremediation?**
K.C. Makris, K.M. Shakya, R. Datta, D. Sarkar, D. Pachanoor

Vetiver grass demonstrates ability to absorb TNT in aqueous media.

5 **Assessment of the impact of increasing concentrations of ozone on photosynthetic components of maize (*Zea mays* L.), a C_4 plant**
L. Leitao, E. Delacôte, P. Dizengremel, D. Le Thiec, J.-P. Biolley

Ozone may induce opposite effects on maize photosynthesis.

Review

9 **Implications of research on endocrine disruption for the environmental risk assessment, regulation and monitoring of chemicals in the European Union**
P. Matthiessen, I. Johnson

The challenges associated with the environmental risk assessment and regulation of endocrine disrupting chemicals are discussed.

Regular Papers

19 **Effect of amendments on the extractability, retention and plant uptake of metals on a sewage-irrigated soil**
B. Paulose, S.P. Datta, R.K. Rattan, P.K. Chhonkar

Liming reduces the phytoavailability of Zn and Ni in metal contaminated soil, while single superphosphate was effective against Cu.

25 **Halogenated organic contaminants and mercury in northern fulmars (*Fulmarus glacialis*): Levels, relationships to dietary descriptors and blood to liver comparison**
L.B. Knudsen, K. Borgå, E.H. Jørgensen, B. van Bavel, M. Schlabach, J. Verreault, G.W. Gabrielsen

Partial dioxin-like toxicity in northern fulmars based on toxic equivalents (TEQs) was well above thresholds for reproductive effects in seabirds.

34 **Dynamic responses of photosystem II in the Namib Desert shrub, *Zygophyllum prismatocarpum*, during and after foliar deposition of limestone dust**
P.D.R. van Heerden, G.H.J. Krüger, M. Kilbourn Louw

Limestone dust deposition reduced photosynthetic capacity in the Namib Desert shrub, Zygophyllum prismatocarpum.

46 **Vegetable and synthetic tannins induce hormesis/toxicity in sea urchin early development and in algal growth**
E. De Nicola, S. Meriç, M. Gallo, M. Iaccarino, C. Della Rocca, G. Lofrano, T. Russo, G. Pagano

Vegetable tannin and synthetic tannins were moderately toxic or displayed hormetic effects in sea urchins and in algae. Re-defining control quality is needed for evaluating hormetic effects.

55 **Effect of physico-chemical pretreatment on the removal efficiency of horizontal subsurface-flow constructed wetlands**
A. Caselles-Osorio, J. Garcia

A physico-chemical pretreatment may help to reduce the risk of clogging of subsurface-flow constructed wetlands.

64 **Nematode communities in contaminated river sediments**
P. Heininger, S. Höss, E. Claus, J. Pelzer, W. Traunspurger

Nematode community structure of river sediments is related to pollution and site structure.

77 **Atmospheric deposition inputs and effects on lichen chemistry and indicator species in the Columbia River Gorge, USA**
M.E. Fenn, L. Geiser, R. Bachman, T.J. Blubaugh, A. Bytnerowicz

Nitrogen, sulfur and acidic deposition threaten natural and cultural resources in the Columbia River Gorge National Scenic Area.

92 **Development and application of long-term sublethal whole sediment tests with *Arenicola marina* and *Corophium volutator* using Ivermectin as the test compound**
Y.T. Allen, J.E. Thain, S. Haworth, J. Barry

Long-term sublethal sediment tests are more sensitive than standard acute tests when assessing the toxic effects of Ivermectin.

doi: 10.1016/S0269–7491(07)00051-6

100 Investigation of organochlorine pesticides (OCPs) in mollusks collected from coastal sites along the Chinese Bohai Sea from 2002 to 2004
Y. Wang, R. Yang, G. Jiang

Concentration of organochlorine pesticides in mollusks collected from coastal sites along the Chinese Bohai Sea didn't change obviously from 2002 to 2004.

107 Occurrence of polybrominated diphenyl ethers (PBDEs) in brown trout bile and liver from Swiss rivers
P.C. Hartmann, P. Burkhardt-Holm, W. Giger

PBDEs with the most abundant BDE-47 were determined in brown trout bile and liver from Swiss rivers.

114 Plant senescence: A mechanism for nutrient release in temperate agricultural wetlands
R. Kröger, M.M. Holland, M.T. Moore, C.M. Cooper

Release of sequestered nutrients during plant senescence.

120 Phytoextraction of excess soil phosphorus
N.C. Sharma, D.L. Starnes, S.V. Sahi

Crop plants such as cucumber, squash and sunflower accumulate phosphorus and thus can be used in the phytoextraction of excess phosphorus from soils.

128 Environmental contaminants in arctic foxes (*Alopex lagopus*) in Svalbard: Relationships with feeding ecology and body condition
E. Fuglei, J.O. Bustnes, H. Hop, T. Mørk, H. Björnfoth, B. van Bavel

High contaminant concentrations in the coastal ecotype of arctic fox may cause toxic health effects due to huge annual cyclic variation in storage and mobilisation of adipose tissue.

139 Impact of electrokinetic remediation on microbial communities within PCP contaminated soil
G. Lear, M.J. Harbottle, G. Sills, C.J. Knowles, K.T. Semple, I.P. Thompson

Electrokinetics negatively impacted soil.

147 Halophyte vegetation influences in salt marsh retention capacity for heavy metals
R. Reboreda, I. Caçador

The type of vegetal cover can affect the overall retention capacity of a salt marsh as well as the functioning of the salt marsh as a sink or source of metals to the estuarine system.

155 A practical demonstration in modelling diclofenac and propranolol river water concentrations using a GIS hydrology model in a rural UK catchment
A.C. Johnson, V. Keller, R.J. Williams, A. Young

A GIS hydrology model was used to predict pharmaceutical concentration hot spots in a rural catchment.

166 Age and seasonal variability of polybrominated diphenyl ethers in free-ranging East Greenland polar bears (*Ursus maritimus*)
R. Dietz, F.F. Rigét, C. Sonne, R.J. Letcher, S. Backus, E.W. Born, M. Kirkegaard, D.C.G. Muir

Mean ΣPBDE concentrations of East Greenland polar bears (Ursus maritimus) were 69.5 ng/g lw (range: 21.7–192 ng/g lw) and showed a seasonal pattern with no relationship with age and sex.

174 Pentachlorophenol association with fulvic acids from recycled wastes
J.C.G. Esteves da Silva, M.C.P.O. Marques

Partition coefficients between pentachlorophenol and fulvic acids extracted form recycled wastes.

180 Effect of drying on the desorption of diuron and terbuthylazine from natural soils
B. Lennartz, X. Louchart

Drying of soil samples increased the binding of herbicidal compounds which is interpreted as a reduction of diffusional mass transfer into and out of the soil organic matter.

188 Congener distribution of polybrominated diphenyl ethers in feral carp (*Cyprinus carpio*) from the Llobregat River, Spain
A. Labandeira, E. Eljarrat, D. Barceló

Differences in PBDE profiles between sediments and fish were observed, suggesting biotransformation in carp.

196 Historical trends in persistent organic pollutants and metals recorded in sediment from Lake Ellasjøen, Bjørnøya, Norwegian Arctic
A. Evenset, G.N. Christensen, J. Carroll, A. Zaborska, U. Berger, D. Herzke, D. Gregor

Levels of Polybrominated diphenyl ethers are increasing in sediment from a Norwegian, Arctic lake.

206 Florida seagrass habitat evaluation: A comparative survey for chemical quality
M.A. Lewis, D.D. Dantin, C.A. Chancy, K.C. Abel, C.G. Lewis

Some anthropogenic contaminants were prevalent in seagrass and their rooted sediments but the biological significance is unknown.

219 Methane and nitrous oxide emission from bovine manure management practices in India
P.K. Gupta, A.K. Jha, S. Koul, P. Sharma, V. Pradhan, V. Gupta, C. Sharma, N. Singh

Emission of CH_4 and N_2O from bovine manure management in India is discussed.

225 Organohalogen compounds in human breast milk from Republic of Buryatia, Russia
O.V. Tsydenova, A. Sudaryanto, N. Kajiwara, T. Kunisue, V.B. Batoev, S. Tanabe

People in the Republic of Buryatia, Russia are exposed to relatively high levels of HCHs, HCB and PCBs.

233 Comparing sediment quality in Spanish littoral areas affected by acute (*Prestige*, 2002) and chronic (Bay of Algeciras) oil spills
C. Morales-Caselles, J. Kalman, I. Riba, T.A. DelValls

Littoral sediments affected by low or moderated but continuous oil spills are more polluted than those affected by accidental oil spills such as the Prestige.

241 Assessment of nitrate pollution in the Grand Morin aquifers (France): Combined use of geostatistics and physically based modeling
N. Flipo, N. Jeannée, M. Poulin, S. Even, E. Ledoux

Combined use of geostatistics and physically based modeling allows assessment of nitrate concentrations in aquifer systems.

257 Residues of organochlorinated pesticides in soils from the Czech Republic
P. Shegunova, J. Klánová, I. Holoubek

Organochlorinated pesticides persist in soils.

262 Accumulation patterns of organochlorines in juveniles of *Arctocephalus australis* found stranded along the coast of Southern Brazil
G. Fillmann, L. Hermanns, T.W. Fileman, J.W. Readman

Analysis of tissues of juveniles of the South American fur seal suggests organochlorines are remobilized during their poor nutritional status, increasing their vulnerability to toxic effects.

268 Organochlorines in harbor seal (*Phoca vitulina*) tissues from the northern Gulf of Alaska
D. Wang, S. Atkinson, A. Hoover-Miller, S.-E. Lee, Q.X. Li

Σ28 PCB congeners, p,p′-DDE, p,p′-DDD and p,p′-DDT, α-, β-, γ-, and δ-HCH and heptachlor in harbor seal tissues from the northern Gulf of Alaska are reported.

281 Priority organic micropollutants in water sources in Flanders and the Netherlands and assessment of removal possibilities with nanofiltration
A. Verliefde, E. Cornelissen, G. Amy, B. Van der Bruggen, H. van Dijk

The article provides a quick and powerful prediction tool for the removal of organic micropollutants by nanofiltration, based on readily available parameter values.

Number 2 2007

SPECIAL ISSUE SECTION: Lichens in a Changing Pollution Environment

291 Lichens in a changing pollution environment: An introduction
O.W. Purvis, M.R.D. Seaward, S. Loppi

293 Further evidence of the effects of global warming on lichens, particularly those with *Trentepohlia* phycobionts
A. Aptroot, C.M. van Herk

Epiphytic and terricolous lichens in Western Europe respond to global warming through their Trentepohlia algae.

299 Diversity and sensitivity of epiphytes to oxides of nitrogen in London
L. Davies, J.W. Bates, J.N.B. Bell, P.W. James, O.W. Purvis

Epiphytic biodiversity in London is related to ambient NOx.

311 Lichen biomonitoring of ammonia emission and nitrogen deposition around a pig stockfarm
L. Frati, S. Santoni, V. Nicolardi, C. Gaggi, G. Brunialti, A. Guttova, S. Gaudino, A. Pati, S.A. Pirintsos, S. Loppi

The diversity of strictly nitrophytic lichen species can be used to monitor and map NH_3 pollution in the Mediterranean area.

317 Is the diversity of epiphytic lichens a reliable indicator of air pollution? A case study from Italy
P. Giordani

Lichen diversity is a reliable tool to monitor atmospheric pollution, but care should be taken in forested areas, where other influencing factors occur.

324 Epiphytic lichens as biomonitors of atmospheric pollution in Slovenian forests
Z. Jeran, T. Mrak, R. Jaćimović, F. Batič, D. Kastelec, R. Mavsar, P. Simončič

No relationship between Hypogymnia element concentrations and foliose epiphytic lichen cover was found.

332 Lichen and bryophyte distribution on oak in London in relation to air pollution and bark acidity
R.S. Larsen, J.N.B. Bell, P.W. James, P.J. Chimonides, F.J. Rumsey, A. Tremper, O.W. Purvis

Transport-related pollutants and bark acidity influence lichen and bryophyte distribution and abundance in London today.

341 Epiphytic lichen community dynamics in deciduous forests around a phosphorus fertiliser factory in Central Lithuania
J. Motiejūnaitė

Biotic factors are important when evaluating dynamics of epiphytic communities in polluted areas.

350 Epiphytic lichens as indicators of environmental quality in Rome
S. Munzi, S. Ravera, G. Caneva

Studies over the past 20 years have characterized the lichen diversity in the city of Rome.

359 Epiphytic lichen growth abnormalities and element concentrations as early indicators of forest decline
T. Otnyukova

Differences in the morphology and element content of Usnea species at two distinct heights of the Abies sibirica canopy provide an early indication of forest decline.

366 Epiphytic macrolichens as indicators of environmental alteration in northern Thailand
W. Saipunkaew, P.A. Wolseley, P.J. Chimonides, K. Boonpragob

Epiphytic lichens in urban and rural areas of northern Thailand can be used as indicators of the levels of pollution associated with increasing population.

375 Response of epiphytic lichen communities to decreasing ammonia air concentrations in a moderately polluted area of The Netherlands
L.B. Sparrius

Lichens can be used to detect both increasing and decreasing ammonia air concentrations.

380 Lichen and moss bags as monitoring devices in urban areas. Part I: Influence of exposure on sample vitality
M. Tretiach, P. Adamo, R. Bargagli, L. Baruffo, L. Carletti, P. Crisafulli, S. Giordano, P. Modenesi, S. Orlando, E. Pittao

The lichen Pseudevernia furfuracea was more resilient than the moss Hypnum cupressiforme in two exposure experiments on trace metal uptake.

392 Lichen and moss bags as monitoring devices in urban areas. Part II: Trace element content in living and dead biomonitors and comparison with synthetic materials
P. Adamo, P. Crisafulli, S. Giordano, V. Minganti, P. Modenesi, F. Monaci, E. Pittao, M. Tretiach, R. Bargagli

Accumulation of trace elements by Hypnum cupressiforme and Pseudevernia furfuracea exposed in bags in urban environment is not influenced by organism vitality.

Review

400 Persistent organic pollutants in human breast milk from Asian countries
S. Tanabe, T. Kunisue

Contamination aspects of POPs in human breast milk from Asian countries were characterized.

Regular Papers

414 Arsenic immobilization in soils amended with drinking-water treatment residuals
D. Sarkar, K.C. Makris, V. Vandanapu, R. Datta

Land-applied drinking-water treatment residuals immobilize arsenic in soils.

420 Hemocyte-specific responses to the peroxidizing herbicide fomesafen in the pond snail *Lymnaea stagnalis* (Gastropoda, Pulmonata)
J. Russo, L. Lefeuvre-Orfila, L. Lagadic

Fomesafen inhibited phagocytosis and the associated oxidative burst, and increased lysosome fragility in L. stagnalis hemocytes.

428 Metal accumulation in the earthworm *Lumbricus rubellus*. Model predictions compared to field data
K. Veltman, M.A.J. Huijbregts, M.G. Vijver, W.J.G.M. Peijnenburg, P.H.F. Hobbelen, J.E. Koolhaas, C.A.M. van Gestel, P.C.J. van Vliet, A. Jan Hendriks

Earthworm metal concentrations are less than linearly related to total soil concentrations and predicted pore water concentrations.

437 Revisiting the plant hyperaccumulation criteria to rare plants and earth abundant elements
C. Branquinho, H.C. Serrano, M.J. Pinto, M.A. Martins-Loução

Revisiting plant hyperaccumulation criteria.

444 Impact of humic substances on the aqueous solubility, uptake and bioaccumulation of platinum, palladium and rhodium in exposure studies with *Dreissena polymorpha*
B. Sures, S. Zimmermann

Precious metal accumulation in Dreissena polymorpha is affected by humic substances.

452 Effect of arbuscular mycorrhizal fungus (*Glomus caledonium*) on the accumulation and metabolism of atrazine in maize (*Zea mays* L.) and atrazine dissipation in soil
H. Huang, S. Zhang, X.-q. Shan, B.-D. Chen, Y.-G. Zhu, J.N.B. Bell

Arbuscular mycorrhizal fungus increases the accumulation and metabolism of atrazine in maize.

458 Biomonitoring of airborne mercury with perennial ryegrass cultures
L. De Temmerman, N. Claeys, E. Roekens, M. Guns

Grass accumulates TGM as a function of the atmospheric concentration and exposure time.

463 Ecotoxicological evaluation of the short term effects of fresh and stabilized textile sludges before application in forest soil restoration
E.V.C. Rosa, T.M. Giuradelli, A.X.R. Corrêa, L.R. Rörig, P.R. Schwingel, C. Resgalla, C.M. Radetski

Short term ecotoxicity evaluation of textile sludge showed that stabilized sludge can be used in the restoration of a non-productive forest soil.

470 Residual concentrations of micropollutants in benthic mussels in the coastal areas of Bohai Sea, North China
W.X. Liu, J.L. Chen, X.M. Lin, Y.S. Fan, S. Tao

Residual Cd contents of mussels at many sites in Bohai Sea exceed national marine biological quality standard and compositions of DDT metabolites indicate recent inputs and potential risk.

478 Prokaryotic life in a potash-polluted marsh with emphasis on N-metabolizing microorganisms
S. Eilmus, C. Rösch, H. Bothe

Characterization of a prokaryotic community of a potash marsh provided information on the occurrence of many unusual prokaryotes and their horizontal distribution.

492 Partitioning and source diagnostics of polycyclic aromatic hydrocarbons in rivers in Tianjin, China
Z. Shi, S. Tao, B. Pan, W.X. Liu, W.R. Shen

Distribution of PAHs among water, suspended solids and sediment was under strong influence of naturally occurring organic carbon.

501 Household hazardous waste disposal to landfill: Using LandSim to model leachate migration
R.J. Slack, J.R. Gronow, D.H. Hall, N. Voulvoulis

Aquatic pollutants linked to the disposal of household hazardous waste in municipal landfills have the potential to exist in soil and groundwater for many years.

Emission of CH_4 and N_2O from bovine manure management in India is discussed.

510 TOL plasmid transfer during bacterial conjugation *in vitro* and rhizoremediation of oil compounds *in vivo*
M.M. Jussila, J. Zhao, L. Suominen, K. Lindström

Horizontal transfer of degradation plasmids in the oil-contaminated rhizosphere reveals the dynamic nature of the intrinsic biodegradation potential.

525 Combined effect of copper and prolonged summer drought on soil Microarthropods in the field
M. Holmstrup, K. Maraldo, P.H. Krogh

Soil contamination with copper does not cause reduced drought tolerance of microarthropods in the field.

534 Interception of residual nitrate from a calcareous alluvial soil profile on the North China Plain by deep-rooted crops: A ^{15}N tracer study
X.T. Ju, Q. Gao, P. Christie, F.S. Zhang

Deep-rooted crops have a greater capacity than shallow-rooted crops to intercept residual nitrate from the subsoil and restrict its movement down to the shallow groundwater.

543 Determination of selected quaternary ammonium compounds by liquid chromatography with mass spectrometry. Part II. Application to sediment and sludge samples in Austria
E. Martínez-Carballo, C. González-Barreiro, A. Sitka, N. Kreuzinger, S. Scharf, O. Gans

The presence of quaternary ammonium surfactants in sediment and sludge samples is reported for the first time in Austria.

548 Bioaccumulation of trace elements in pelagic fish from the Western Indian Ocean
J. Kojadinovic, M. Potier, M. Le Corre, R.P. Cosson, P. Bustamante

Metal bioaccumulation was quantified in four species of pelagic fish.

567 Landscape ecology of the Guanting Reservoir, Beijing, China: Multivariate and geostatistical analyses of metals in soils
W. Luo, T. Wang, Y. Lu, J.P. Giesy, Y. Shi, Y. Zheng, Y. Xing, G. Wu

Concentrations of metals, sources and spatial distributions in soils around Guanting Reservoir were determined.

Number 3 2007

SPECIAL ISSUE: Critical Levels for Ozone Effects on Vegetation: Further Applying and Developing the Flux Concept

Guest Editorial

577 Preface to the special issue of the Obergurgl workshop on ozone
G. Wieser, M. Tausz

578 Robustness of modelled ozone exposures and doses
J.-P. Tuovinen, D. Simpson, L. Emberson, M. Ashmore, G. Gerosa

The $AF_{st}Y$ index is less sensitive to uncertainties than AOTX, but depends strongly on the stomatal conductance model.

587 Promoting the O_3 flux concept for European forest trees
R. Matyssek, A. Bytnerowicz, P.-E. Karlsson, E. Paoletti, M. Sanz, M. Schaub, G. Wieser

Flux-based O_3 risk assessment of forest trees copes with variable moisture and light effects and requires the effective O_3 dose to account for the trees' O_3 defence and tolerance.

608 Risk assessments for forest trees: The performance of the ozone flux versus the AOT concepts
P.E. Karlsson, S. Braun, M. Broadmeadow, S. Elvira, L. Emberson, B.S. Gimeno, D. Le Thiec, K. Novak, E. Oksanen, M. Schaub, J. Uddling, M. Wilkinson

Ozone stomatal flux based indices were superior, as compared to AOT40, for explaining biomass reductions and leaf visible injury

617 Simulation of stomatal conductance for Aleppo pine to estimate its ozone uptake
S. Elvira, R. Alonso, B.S. Gimeno

Current EMEP stomatal uptake module needs to be re-parameterised for Mediterranean tree species.

624 Shoot growth of mature *Fagus sylvatica* and *Picea abies* in relation to ozone
S. Braun, C. Schindler, B. Rihm, W. Flückiger

Epidemiological assessment of shoot growth suggests an ozone sensitivity of mature beech which is similar to seedlings.

629 Exemplifying whole-plant ozone uptake in adult forest trees of contrasting species and site conditions
A.J. Nunn, G. Wieser, U. Metzger, M. Löw, P. Wipfler, K.-H. Häberle, R. Matyssek

Sap flow-based assessment of whole-tree O_3 uptake reflects similar responsiveness of canopy conductance and O_3 uptake across contrasting tree species and site conditions.

640 Comparison of calculated and measured foliar O_3 flux in crop and forest species
N.E. Grulke, E. Paoletti, R.L. Heath

Using a new system to concurrently measure H_2O, O_3, and CO_2 flux, the conventional method of calculating O_3 flux generally overestimated direct measures by 25–50%.

648 Measuring, modelling and testing ozone exposure, flux and effects on vegetation in southern European conditions—What does not work? A review from Italy
M. Ferretti, M. Fagnano, T. Amoriello, M. Badiani, A. Ballarin-Denti, A. Buffoni, F. Bussotti, A. Castagna, S. Cieslik, A. Costantini, A. De Marco, G. Gerosa, G. Lorenzini, F. Manes, G. Merola, C. Nali, E. Paoletti, B. Petriccione, S. Racalbuto, G. Rana, A. Ranieri, A. Tagliaferri, G. Vialetto, M. Vitale

While the flux-based approach is scientifically sounder, a more practical concentration-based approach is still necessary for routine monitoring.

659 Modelling stomatal ozone flux and deposition to grassland communities across Europe
M.R. Ashmore, P. Büker, L.D. Emberson, A.C. Terry, S. Toet

Modelled stomatal flux of ozone to productive grasslands in Europe shows different spatial and temporal variation to AOT40, and is modified by management and soil water status.

671 High-resolution modelling of AOT40 and stomatal ozone uptake in wheat and grassland: A comparison between 2000 and the hot summer of 2003 in Switzerland
F. Keller, S. Bassin, C. Ammann, J. Fuhrer

A national-scale assessment based on stomatal ozone flux, but not on AOT40, revealed similar risks for crops in years with different climates and ozone levels.

678 Factors affecting the ozone sensitivity of temperate European grasslands: an overview
S. Bassin, M. Volk, J. Fuhrer

An overview of experimentally induced ozone effects suggests that temperate grasslands could be separated into broad classes of ozone sensitivity based on physiological and ecological principles.

692 Stomatal conductance of semi-natural Mediterranean grasslands: Implications for the development of ozone critical levels
R. Alonso, V. Bermejo, J. Sanz, B. Valls, S. Elvira, B.S. Gimeno

The stomatal conductance model incorporated within the EMEP DO_3SE deposition module needs to be re-parameterised for Mediterranean semi-natural vegetation.

699 Derivation of ozone flux-yield relationships for lettuce: A key horticultural crop
E. Goumenaki, I.G. Fernandez, A. Papanikolaou, D. Papadopoulou, C. Askianakis, G. Kouvarakis, J. Barnes

Establishment of ozone flux-yield relationships for a commercially-important horticultural crop grown widely in the Mediterranean.

707 Ozone stress and antioxidant substances in *Trifolium repens* and *Centaurea jacea* leaves
J.F. Severino, K. Stich, G. Soja

Low leaf ascorbic acid levels are a main cause for visible ozone injuries in Trifolium and Centaurea.

715 A comparison of two different approaches for mapping potential ozone damage to vegetation. A model study
D. Simpson, M.R. Ashmore, L. Emberson, J.-P. Tuovinen

Model calculations of AOT40 and AFstY show very different spatial variations in the risks of ozone damage to vegetation.

726 Comparison of different stomatal conductance algorithms for ozone flux modelling
P. Büker, L.D. Emberson, M.R. Ashmore, H.M. Cambridge, C.M.J. Jacobs, W.J. Massman, J. Müller, N. Nikolov, K. Novak, E. Oksanen, M. Schaub, D. de la Torre

A comparison of an empirical and a semi-mechanistic stomatal conductance algorithm for European crop and forest species reveals a similar predictive capability.

736 Identifying ozone-sensitive communities of (semi-)natural vegetation suitable for mapping exceedance of critical levels
G. Mills, F. Hayes, M.L.M. Jones, S. Cinderby

Grassland communities such as alpine and sub-alpine grasslands have the highest potential sensitivity ozone, based on the responses of their component species.

744 Predicting community sensitivity to ozone, using Ellenberg Indicator values
M.L.M. Jones, F. Hayes, G. Mills, T.H. Sparks, J. Fuhrer

The ozone sensitivity of individual species can be predicted from Ellenberg Light and Salinity Indicator values and extrapolated to the community level.

754 Meta-analysis of the relative sensitivity of semi-natural vegetation species to ozone
F. Hayes, M.L.M. Jones, G. Mills, M. Ashmore

Meta-analysis of the relative sensitivity of semi-natural vegetation species to ozone showed some relationships with physiological and ecological characteristics.

763 Implications of climate change for the stomatal flux of ozone: A case study for winter wheat
H. Harmens, G. Mills, L.D. Emberson, M.R. Ashmore

In a future climate the stomatal flux of ozone is likely to be reduced across Europe despite increasing tropospheric background ozone concentrations.

I Contents of Volume 146